



UNIVERSAL BEER AGAR (7574)

Intended Use

Universal Beer Agar is used for the cultivation of bacteria and yeasts encountered in the brewing industry.

Product Summary and Explanation

Universal Beer Agar was developed by Kozulis and Page, and used as a basal medium supplemented with beer.¹ The developers compared this formula with other media commonly used in breweries for detecting microbial contamination.² The characteristics of Universal Beer Agar are closer to natural environmental conditions found in a typical brewery. Universal Beer Agar supports the growth of more lactic acid bacteria and yields larger colonies in a shorter time than traditional brewer's media. Supplementing this medium with beer has produced a selective environment for organisms that are adapted to existent conditions in the brewery. The presence of hop constituents and alcohol inhibits growth of many airborne microorganisms not adapted to this environment.³

Universal Beer Agar supports growth of *Lactobacillus*, *Pediococcus*, *Acetobacter*, and yeasts that are known beer contaminants. Universal Beer Agar is abbreviated as UBA.

Principles of the Procedure

Tomato Juice Solids is a source of carbon, protein, and nutrients. Yeast Extract provides trace elements, vitamins, and amino acids. Dextrose supplies additional carbon. Dipotassium Phosphate and Monopotassium Phosphate are the buffers. Magnesium Sulfate, Ferrous Sulfate, and Manganese Sulfate are sources of ions that stimulate metabolism. Sodium Chloride maintains the osmotic balance of the medium. Peptonized Milk contains lactose as an energy source. Agar is the solidifying agent.

Formula / Liter

Tomato Juice Solids	7 g
Yeast Extract.....	10 g
Dextrose.....	10 g
Dipotassium Phosphate	0.5 g
Monopotassium Phosphate	0.5 g
Magnesium Sulfate	0.125 g
Sodium Chloride	0.01 g
Ferrous Sulfate	0.01 g
Manganese Sulfate	0.01 g
Peptonized Milk.....	15 g
Agar	12 g

Final pH: 6.3 ± 0.2 at 25°C

Formula may be adjusted and/or supplemented as required to meet performance specifications.

Precaution

1. For Laboratory Use.

Directions

1. Suspend 55 g of the medium in 750 mL of purified water.
2. Heat with frequent agitation and boil for one minute to completely dissolve the medium.
3. While medium is hot, add and mix 250 mL of beer without degassing.
4. Autoclave at 121°C for 10 minutes.

Quality Control Specifications

Dehydrated Appearance: Powder is homogeneous, free flowing, and light brown.

Prepared Appearance: Prepared medium is trace to slightly hazy and light to dark amber.

Expected Cultural Response: Cultural response on Universal Beer Agar incubated aerobically at 25 - 30°C and examined for growth after 18 - 72 hours.

Microorganism	Approx. Inoculum (CFU)	Expected Results
<i>Escherichia coli</i> ATCC® 25922	10 - 300	Growth
<i>Lactobacillus casei</i> ATCC® 393	10 - 300	Growth
<i>Lactobacillus fermentum</i> ATCC® 9338	10 - 300	Growth
<i>Saccharomyces cerevisiae</i> ATCC® 9763	10 - 300	Growth

The organisms listed are the minimum that should be used for quality control testing.

Test Procedure

Refer to appropriate references for specific procedures.

Results

Refer to appropriate references and procedures for results.

Storage

Store sealed bottle containing the dehydrated medium at 2 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light by keeping container tightly closed.

Expiration

Refer to expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

Limitation of the Procedure

Due to varying nutritional requirements, some strains may be encountered that grow poorly or fail to grow on this medium.

Packaging

Universal Beer Agar	Code No.	7574A	500 g
		7574B	2 kg
		7574C	10 kg

References

1. **Kozulis, J. A., and H. E. Page.** 1968. A new universal beer agar medium for the enumeration of wort and beer microorganisms. Proc. Am. Soc. Brew. Chem. **19**:52-58.
2. **Murphy, D. T., and L. T. Saletan.** 1970. Use of microbiological media in the brewery. Tech. Q. Master Brew. Assoc. Am. **7**:182-187.
3. **MacFaddin, J. D.** 1985. Media for isolation-cultivation-identification-maintenance medical bacteria, vol. 1. p. 819-820. Williams & Wilkins, Baltimore, MD.

Technical Information

Contact Acumedia Manufacturers, Inc. for Technical Service or questions involving dehydrated culture media preparation or performance at (517)372-9200 or fax us at (517)372-2006.