

LETHEEN AGAR WITH TWEEN (7710)

Intended Use

Letheen Agar with Tween is used for the testing of quaternary ammonium compounds for antimicrobial activity.

Product Summary and Explanation

In 1948, Weber and Black described the value of a highly nutritional solid medium containing neutralizing agents for quaternary ammonium compounds in sanitizers.¹ The addition of Lecithin and Tween 80 (Polysorbate 80) to Tryptone Glucose Extract (TGE) Agar resulted in a medium that effectively neutralizes quaternary ammonium compounds while testing germicidal activity. Letheen Agar with Tween is a modification of TGE Agar, with the addition of Lecithin and Tween 80 (Polysorbate 80).

Letheen Agar with Tween is specified for use by the American Society for Testing Materials (ASTM) in Standard Test Method for Preservatives in Water-Containing Cosmetics.² Total neutralization of disinfectants is critical. Disinfectant residues can result in a false negative (no-growth) test.

Letheen Agar with Tween has incorporated Tween 80 (Polysorbate 80) into the medium, this is the only difference from Letheen Agar Base.

Principles of the Procedure

Enzymatic Digest of Casein and Beef Extract provide nitrogen, carbon, vitamins, and minerals in Letheen Agar with Tween. Dextrose is the fermentable carbohydrate. Lecithin neutralizes quaternary ammonium compounds and Tween 80 (Polysorbate 80) neutralizes phenols, hexachlorophene, formalin, and with Lecithin, ethanol.^{3,4,5,6} Agar is the solidifying agent.

Formula / Liter

Enzymatic Digest of Casein	5 g
Dextrose.....	1 g
Beef Extract	3 g
Lecithin.....	1 g
Tween 80 (Polysorbate 80).....	7 g
Agar	15 g

Final pH: 7.0 ± 0.2 at 25°C

Formula may be adjusted and/or supplemented as required to meet performance specifications.

Precaution

1. For Laboratory Use.

Directions

1. Suspend 32 g of the medium in one liter of purified water.
2. Heat with frequent agitation and boil for one minute to completely dissolve the medium.
3. Autoclave at 121°C for 15 minutes.

Quality Control Specifications

Dehydrated Appearance: Powder is homogeneous, lumpy, and tan.

Prepared Appearance: Prepared medium is light to medium amber, and trace to slightly hazy.

Expected Cultural Response: Cultural response on Lethen Agar with Tween incubated aerobically at 35 ± 2°C and examined for growth after 18 - 24 hours.

Microorganism	Approx. Inoculum (CFU)	Response
<i>Enterococcus faecalis</i> ATCC® 29212	10 - 300	Good growth
<i>Escherichia coli</i> ATCC® 25922	10 - 300	Good growth
<i>Pseudomonas aeruginosa</i> ATCC® 27853	10 - 300	Good growth
<i>Salmonella typhimurium</i> ATCC® 14028	10 - 300	Good growth
<i>Staphylococcus aureus</i> ATCC® 25923	10 - 300	Good growth

The organisms listed are the minimum that should be used for quality control testing.

Test Procedure

Lethen Agar with Tween is used in a variety of procedures. Consult appropriate references for complete information.^{4,7}

Results

Refer to appropriate references and procedures for results.

Storage

Store sealed bottle containing the dehydrated medium at 2 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light by keeping container tightly closed.

Expiration

Refer to expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

Limitation of the Procedure

Due to nutritional variation, some strains may be encountered that grow poorly or fail to grow on this medium.

Packaging

Lethen Agar with Tween	Code No.	7710A	500 g
		7710B	2 kg
		7710C	10 kg

References

1. **Weber, G. R., and L. A. Black.** 1948. Relative efficiency of quaternary inhibitors. Soap and Sanit. Chem. **24**:134-139.
2. **American Society for Testing Materials.** 1991. Standard test method for preservatives in water-containing cosmetics, E 640-78. Annual Book of ASTM Standards, Philadelphia, PA.
3. **Quisno, R., I. W. Gibby, and M. J. Foter.** 1946. A neutralizing medium for evaluating the germicidal potency of the quaternary ammonium salts. Am. J. Pharm. **118**:320-323.
4. **Erlanson, A. L., Jr., and C. A. Lawrence.** 1953. Inactivating medium for hexachlorophene (G-11) types of compounds and some substituted phenolic disinfectants. Science. **118**:274-276.
5. **Brummer, B.** 1976. Influence of possible disinfectant transfer on *Staphylococcus aureus* plate counts after contact sampling. Appl. Environ. Microbiol. **32**:80-84.
6. **Favero (chm.).** 1967. Microbiological sampling of surfaces-a state of the art report. Biological Contamination Control Committee, American Association for Contamination Control.
7. **Association of Official Analytical Chemists.** 1995. Official methods of analysis, 16th ed. Association of Official Analytical Chemists, Washington, D.C.

Technical Information

Contact Acumedia Manufacturers, Inc. for Technical Service or questions involving dehydrated culture media preparation or performance at (517)372-9200 or fax us at (517)372-2006.